

(so as a ring  $H_{dR}^*(\mathbb{C}P^n; \mathbb{R}) \cong \mathbb{R}[t]/(t^{n+1}=0)$  polynomial ring up to degree  $n$ .)

### 3. cptly supp de Rham cohomology

This is designed to deal with non-cpt wfd  $M$ .

$\Omega_c^k(M; \mathbb{R}) = \{ \text{cpt supp } k\text{-form on } M \}$  ← This is closed under exterior derivative

KEY observation: it could be  $d: \underset{\uparrow \Omega_c^k(M)}{\alpha} \longrightarrow \underset{\uparrow \Omega_c^{k+1}(M)}{d\alpha}$

BUT  $\alpha$  is not cptly supp. e.g.  $f \equiv 1 \in \Omega_c^0(\mathbb{R}^n)$ .

Define  $H_c^k(M; \mathbb{R}) := \frac{\ker \{ d_k: \Omega_c^k(M; \mathbb{R}) \rightarrow \Omega_c^{k+1}(M; \mathbb{R}) \}}{\text{Im} \{ d_{k-1}: \Omega_c^{k-1}(M; \mathbb{R}) \rightarrow \Omega_c^k(M; \mathbb{R}) \}}$

e.g. compute  $H_c^*(\mathbb{R}^1; \mathbb{R})$

$$H_c^0(\mathbb{R}^1; \mathbb{R}) = \frac{\ker \{d_0: \Omega_c^0(\mathbb{R}^1; \mathbb{R}) \rightarrow \Omega_c^1(\mathbb{R}^1; \mathbb{R})\}}{0}$$

$$= \frac{0}{0} = 0.$$



v/c  $\Omega_c^0(\mathbb{R}^1; \mathbb{R}) \rightarrow 0$   
 $(df=0 \Rightarrow f$  is constant, and it has to be 0 in order to be cply supp)

This is in a sharp contrast to  $H_{loc}^0(\mathbb{R}^1; \mathbb{R}) (\cong \mathbb{R})$ .  
 In general, when  $M$  is non-cpt,  $H_c^0(M; \mathbb{R}) = 0$ .

$$H_c^1(\mathbb{R}^1; \mathbb{R}) = \frac{\ker(d_1: \Omega_c^1(\mathbb{R}^1; \mathbb{R}) \rightarrow 0)}{\text{Im}(d_0: \Omega_c^0(\mathbb{R}^1; \mathbb{R}) \rightarrow \Omega_c^1(\mathbb{R}^1; \mathbb{R}))}$$

$$= \frac{\Omega_c^1(\mathbb{R}^1; \mathbb{R})}{\text{Im}(d_0)} = \frac{\{f(x)dx \mid f \in C_c(\mathbb{R})\}}{\{dg \mid g \in C_c(\mathbb{R})\}}$$

Naive: For any  $f(x)dx$ , consider  $g(x) = \int_{-\infty}^x f(t)dt$  (then  $dg = f(x)dx$ )  
 but  $g$  is not nec inside  $\Omega_c^0(\mathbb{R}^1; \mathbb{R})$ .



(This in fact indicates that  $H^1(\mathbb{R}^1; \mathbb{R}) = 0$ .)

Consider  $S: \Omega_c^1(\mathbb{R}^1; \mathbb{R}) \rightarrow \mathbb{R}$  by  $f(x)dx \mapsto \int_{\mathbb{R}} f(x)dx < \infty$ .

Then  $\ker(S) = \{f(x)dx \mid \int_{\mathbb{R}} f(x)dx = 0\}$ .

By construction above, consider  $g(x) = \int_{-\infty}^x f(t)dt$  for any  $f(x)dx$  in  $\ker(S)$  and  $g \in \text{Im}(d_0)$ . ( $\Rightarrow \ker(S) = \text{im}(d_0)$ ).

$\Rightarrow H_c^1(\mathbb{R}^1; \mathbb{R}) \cong \mathbb{R}$  so  $\dim H_c^1(\mathbb{R}^1; \mathbb{R}) = 1$ .

Recall  $H_c^1(\mathbb{R}^1; \mathbb{R}) = 0$ .

- From  $H_c^0(\mathbb{R}^1; \mathbb{R})$ , we obtain a general result:

$$H_c^0(M; \mathbb{R}) = \mathbb{R}^{\# \text{cpt connected components of } M}.$$

- From  $H_c^0(\mathbb{R}^1; \mathbb{R})$ , we know that  $H_c^*(M; \mathbb{R})$  is not an invariant up to homotopy equivalence (b/c  $\mathbb{R}^1 \cong \{pt\}$  but  $H_c^0(\{pt\}; \mathbb{R}) = \mathbb{R}$ ).

-  $f: \mathbb{R} \rightarrow \mathbb{R}$  by  $f(x) \equiv 0$ , then  $f^*$ :  $\Omega^1(\mathbb{R}; \mathbb{R}) \rightarrow$   
for the pullback  
 $\Omega^1(\mathbb{R}; \mathbb{R})$ ,  $f^*(0) = \Omega^1(\mathbb{R}; \mathbb{R})$  (much bigger than  $\Omega_c^1(\mathbb{R}; \mathbb{R})$ ).

so  $f^*$  does not pullback well in terms of cpt supp forms.

To fix this, one usually consider two variants:

① assume  $f$  is proper (preimage of cpt under  $f$  is cpt).

② instead of  $f^*$ , consider "push-forward  $f_*$ ".

For ①,  $f: N \rightarrow M$  and proper then similarly to the standard case, we have  $f^*: H_c^k(M; \mathbb{R}) \rightarrow H_c^k(N; \mathbb{R})$ .

For ②, only works for special cases:

e.g. If  $N \subset M$  and  $f: N \rightarrow M$  is the inclusion, then define

$f_* (\theta) \in \Sigma_c^k(M; \mathbb{R})$  = extension by zero of  $\theta \in \Sigma_c^k(N; \mathbb{R})$ .

e.g. If  $\begin{matrix} E \\ \downarrow \pi \\ M \end{matrix}$  a vector bundle.  $\theta \in \Sigma_c^k(E)$ , then define

$$(\pi_* (\theta)) (\varphi) := \int_{\pi^{-1}(\varphi)} \theta \in \Sigma_c^k \text{-rank of } \begin{matrix} E \\ \downarrow \\ M \end{matrix} \rightarrow \text{reference Thom class}$$

← called integration along fibers

(DIY) For  $H_c^*(M; \mathbb{R})$ , we also have a MV-seq (but with opposite direction):

$$\dots \rightarrow H_c^k(U \cap V; \mathbb{R}) \rightarrow H_c^k(U; \mathbb{R}) \oplus H_c^k(V; \mathbb{R}) \rightarrow H_c^k(M; \mathbb{R}) \rightarrow \dots$$

Then (Poincaré duality) Let  $M$  be an oriented manifold. Then

$$H_{dR}^*(M; \mathbb{R}) \cong \left( H_c^{\dim M - *}(M; \mathbb{R}) \right)^* \leftarrow \begin{matrix} \text{without} \\ \text{1/d} \end{matrix} \leftarrow \text{dual of a vector space}$$

This isomorphism is explicitly given by  $[\theta] \mapsto$  a linear map PD  $([\theta])$

defined by  $PD([0]) ([0]) := \int_M \theta \lrcorner \sigma$   $\leftarrow$  this is in top degree.

Remark well-definedness of this def.  $\theta + dz$ , then  $\int_M (\theta + dz) \lrcorner \sigma = \int_M \theta \lrcorner \sigma + \int_M d(z \lrcorner \sigma)$   $\leftarrow$   $\int_M d(z \lrcorner \sigma) = 0$  by Stokes' Thm.  $\leftarrow$   $\sigma$  is closed

$= \int_M \theta \lrcorner \sigma.$

e.g. Let  $M = \mathbb{R}^n$ , then

$$H_c^k(\mathbb{R}^n; \mathbb{R}) = (H_{dR}^{n-k}(\mathbb{R}^n; \mathbb{R}))^* = \begin{cases} \mathbb{R} & k = n \\ 0 & k \neq n \end{cases}$$

e.g. If  $M^n$  is connected, non-cpt, orientable

$$H_c^n(M; \mathbb{R}) \simeq (H_{dR}^0(M; \mathbb{R}))^* = \mathbb{R}$$

(NEW)  $H_{dR}^n(M; \mathbb{R}) \simeq (H_c^0(M; \mathbb{R}))^* = 0$

Recall in Lecture 5, we defined and calculated  $H_{dR}^k(M; \mathbb{R})$  when  $M$  is cpt

In class, the lecturing around here is not correct.

For non-orientable manifolds, the Poincaré duality stated above does not apply. I was planning to emphasize the non-trivial fact in **\*\*Rmk\*\*** right below,

where the proof needs to use a fact in algebraic topology:

any non-orientable manifold a double-cover that is orientable.

e.g.  $M$  cpt, non-orientable  $\Rightarrow H_c^n(M; \mathbb{R}) = H_{dR}^n(M; \mathbb{R}) = 0$

Rmk  $M$  non-cpt non-orientable  $\Rightarrow H_c^n(M; \mathbb{R}) = 0$  (see proof in Lee's book Thm 17.34)

(NEW) *We can't apply Poincaré duality as stated above (which works for orientable cases).*

e.g. if  $M$  is closed, then  $H_c^*(M; \mathbb{R}) \cong H_{dR}^*(M; \mathbb{R})$ .

$\Rightarrow M^3$ , then  $H_{dR}^0(M; \mathbb{R}) \cong H_{dR}^3(M; \mathbb{R})$  (\*)

$H_{dR}^1(M; \mathbb{R}) \cong H_{dR}^2(M; \mathbb{R})$ .

$\Rightarrow \chi(M) = \sum_{k=0}^3 (-1)^k b_k(M; \mathbb{R}) = \dim H^0 - \dim H^1 + \dim H^2 - \dim H^3 = 0$ .

(In general, any  $M^{\text{odd}}$  has  $\chi(M) = 0$ ).

Rmk Every odd-dim closed mfd is orientable. (by (\*))

Rmk We will prove Poincaré duality in next section.

To end this section, let us demonstrate an application of  $H_c^*(M; \mathbb{R})$

For  $f: N^n \rightarrow M^n$  proper, consider  $f^*: H_c^n(M; \mathbb{R}) \rightarrow H_c^n(N; \mathbb{R})$   
orientable, connected without b/d.  $\mathbb{R}$   $\mathbb{R}$

Then fix any generators  $\alpha \in H_c^n(M; \mathbb{R})$ , we have

$$\int_N f^* \alpha = \lambda \cdot \int_M \alpha \quad \text{defined by } \int_N \alpha \text{ for any rep } \alpha \text{ of class } \alpha \implies \text{define } \deg(f) = \lambda.$$


degree of  $f$

The degree  $\deg(f)$  is independent of the choice of the generator.

FACT (proved in next lecture)  $\deg(f) \in \mathbb{Z}$ .

Here are trivial observations directly from def.

- If  $f: N \rightarrow M$  is not surjective, then  $\deg(f) = 0$ .





- If  $f: M \rightarrow M$  is the identity map, then  $\deg(f) = 1$ .

-  $L \xrightarrow{f} N \xrightarrow{g} M \Rightarrow \deg(g \circ f) = \deg(f) \deg(g)$ .

$f, g: N \rightarrow M$   
proper homotopic  
then  
 $\deg(f) = \deg(g)$ .


- If  $f: M \rightarrow M$  differs, then  $\forall \alpha \in H_c^n(M; \mathbb{R})$ , we have

$$\int_M f^* \alpha = \pm \int_M \alpha \Rightarrow \deg(f) = \begin{cases} 1 & \text{orientation preserving} \\ -1 & \text{orientation reversing} \end{cases}$$

↑  
change variable

b/c  $\deg(f) \cdot \deg(f^{-1}) = \deg(\text{id}) = 1$   
 $\Rightarrow \deg(f) = \pm 1$

e.g.  $f: S^1 \rightarrow S^1$



$p \rightarrow -p \quad (x, y) \mapsto (-x, -y)$

then take a closed 1-form  $\theta = -y dx + x dy$  for  $(x, y) \in S^1$  (i.e.  $x^2 + y^2 = 1$ ).

$$\int_{S^1} f^* \theta = \int_{S^1} \theta \quad \text{b/c both } x, y \text{ change sign} \Rightarrow \deg(f) = 1$$

One can imagine, for  $S^2$ , the "antipodal" map  $p \rightarrow -p$  will have  $\deg(f) = -1$ .

In general,  $\deg(f: S^n \rightarrow S^n) = \begin{cases} +1 & \text{if } n \text{ is odd} \\ -1 & \text{if } n \text{ is even} \end{cases}$  (antipodal)

$\Rightarrow$  On  $S^{2n}$ , there does not exist any nowhere vanishing vector fields.  
*general hairy ball theorem*

Pf. Set  $S^{2n} \subset \mathbb{R}^{2n+1}$  as the standard sphere, then vector field  $X$  at pt  $p \in S^{2n}$  lies in the tangent plane  $T_p S^{2n}$ , which is orthogonal to  $p \in \mathbb{R}^{2n+1}$ . In particular,  $X(p), p$  are always linearly indep in  $\mathbb{R}^{2n+1}$ .

Suppose  $X(p) \neq 0 \forall p$ , then consider a htp  $\leftarrow$  a continuous map param by  $t$ .  
 $\leftarrow$  since  $X(p) \neq 0 \forall p$ .

$$\cos(\pi t) p + \sin(\pi t) X(p) \text{ for } t \in [0, 1]$$

Then maps  $f_t: p \mapsto \cos(\pi t) p + \sin(\pi t) X(p)$  is a htp from

$$f_0 = \mathbb{1} \text{ to } f_1 = \text{antipodal map} \Rightarrow f_0^* \alpha = f_1^* \alpha \Rightarrow \deg(f_0) = \deg(f_1) \rightarrow \leftarrow$$

Prop  $M^n$  cpt mfd with b/d  $\partial M$

$X^{n-1}$  cpt orientable mfd

If  $f: \partial M \rightarrow X^{n-1}$  can be extended to  $g: M \rightarrow X^{n-1}$ ,

then  $\deg(f) = 0$

(How to apply: suppose  $X^{n-1} = \partial M$  for some  $M$ , say  $M = B^n$   $n$ -dim ball, then  $X^{n-1} = \partial M = S^{n-1}$ . Under the hypothesis above, such  $f$  can not be homotopic to either  $\mathbb{1}$  or antipodal map!)

Pf of prop. Fix a volume form  $\Omega$  on  $X^{n-1}$  s.t.  $\int_X \Omega = 1$

Then  $\deg(f) = \deg(f) \int_X \Omega = \int_{\partial M} f^* \Omega$ .

$$\begin{array}{ccc} \partial M & \xrightarrow{i} & M \xrightarrow{g} X^{n-1} \\ & \searrow & \uparrow \\ & & f = g \circ i \end{array} \Rightarrow \int_{\partial M} f^* \Omega = \int_{\partial M} i^*(g^* \Omega) = \int_M d(g^* \Omega) = \int_M g^* d\Omega = 0$$

## Extension reading topics

- de Rham cohomology groups on a Lie group

← Chevalley-Eilenberg's original paper 1948

- Thom class (integration over fiber)

← Bott-Tu's book, Chapter I

- Poincaré duality of manifolds (intersection theory; basic).

↑  
notes from Nico Iaescu (Notre Dame)