## Fredholm analysis

## O. Rewite the past

$$\mathcal{E} = \mathcal{O}(10) \times \mathcal{D}^{el}(\mathcal{E}, u^*TM) \qquad \text{approbe} \qquad \mathcal{E} = \mathcal{O}(10) \times W^{k-1,p}(\mathcal{D}^{el}(\mathcal{E}, u^*TM))$$

$$\mathcal{B} = \mathcal{O}(\mathcal{E}, M) \qquad \text{Sobilev space} \qquad \mathcal{B} = W^{k,p}(\mathcal{E}, M)$$

Here one should view

· 
$$W^{k,p}(\Sigma,M) = \overline{C^{\infty}(\Sigma,M)}^{k,l}(\Sigma^{0,l}(\Sigma,U^{*TM}))$$

· WER (E,M) is defined locally from WER (D', C").

Thenfor, the last theorem in SFT-3 can be globally stated:

Thu  $U: (\overline{\Sigma}_j) \to (M, \overline{J})$   $\overline{J}$ -hol and  $\overline{J}$  is  $C^{\kappa}$ , when if  $U \in W^{k,p}(\underline{\Sigma},M)$  for p>2, when  $U \in W^{k,p}(\underline{\Sigma},M)$ . In particular, if  $\overline{J}$  is smooth, when  $U \in C^{\infty}(\underline{\Sigma},M)$ .

Often one simply the whaten by denoting Du (=(\$\overline{c}\_{2})(")).

## 1. Basic Fredholm

= n-w

A bounded linear operator D: X -> Y 17 called a Fredholm operator

Banach space (over 12 m C)

if (1) im(D) is closed in Y; (2) dim (Ker D) is finite: (3) dim (1/1m01)
is also finite.

= ind(D) = dim ker D - dim coker D called Fredholm index

Ex X= R" Y= Rm, then any linear map D: X > Y is Fredholm.

Note that this ind (D) is in fact independent of D. In au-dim/

Question: How to venify that DiX > Y is Fredholm?

Lemma (very useful) D: X > Y bld linear operator i and dim (Kens) case that depend on D, K K: X -> Z cpt operator i toker, one usually

then D has a closed image and dim(terD) = The two conclusions also supp D to be called sensitively

=> starting from D in the conclusion, assume dim KerD = n.

then I Ku: X > IR" (coming from projection to the complement
of KerD) st. Ko | KerD cx: KerD ~ IR"

comes from
ttahu-Bound The

 $\Rightarrow \quad X \to \mathcal{T} \otimes \mathbb{R}^n \qquad x \mapsto (Dx, kx) \text{ is injective}$  and has closed image.  $\left(so\ \exists\ \text{bijective}:\ X \to \text{image of }(D, k_0)\right)$ 

Barach iso thus

The inverse is a bounded operator, i.e. I (>0

(from spen wappy thus)

s.t.  $||x||_X \leq C \left( ||Dx||_Y + ||K_0x||_{\mathbb{R}^n} \right)$  ind of p. therefor, if  $p: X \to Y$  satisfying  $||p|| = \frac{1}{C}$  then

11x11x - C 11px1/= ( (11p+p)x1/4+11 bx1/4+11/8x1/18x)- C 11px1/4

pxII = ripil lix(1x

By Leanne above. Dep has a closed image and dim (Ker (D+P)) cas.

By a dual congrument for D (D\*: Y\* -> X\*), we get a simular conclusion:

= \$2 > 0 5.4. & p 1(p)/< 5,

D closed image + dim (coker D) < a >> D+p closed image + dim (coker D) < coo.

Prof. If D: X-y is a Fredlulm operator, then DESO 5.t. & p: X-y with liplics, we have D+P is also Fredholm.

Exercise: in Prop. one can also prove ind (D+p) = ind (D).

Ruk (easy to verify):

- . D: X -x, T: Y -> Z Fredholm => i'nd (TD)=ind (D)+ind (T)
- · D: X > Fredholm => ind (D\*) = ind (D).

Set  $Fred(X,Y):=\{D:X\to Y \text{ bounded } | D \text{ is Fredholm}\}$ One can associate a top

Con If  $\delta: [0,1] \longrightarrow Fred(X, f)$  is a continuous parch of Fredholm operator, then ind (Y(0)) = ind(Y(0)).

## Z Fredholm waps

f: X -> Y a Smooth map and it's called a Fredholm map Banach unfol Cluely modelled by Banach space) if  $\forall x \in X$ ,  $df(x): T_{*}X \rightarrow T_{fin}Y$  is a Fredholm operator.

Important! Assume X is path connected, then x x' = df(x) ~ df(x')

=> (vd(df(x)) is independent of x & X

= denote ind (f) (= ind (df(x)) +x +x).

Recall that in smooth unfol , f: M -> N and concept ingular value! The same concept in Banach unfol setting.

•  $y \in Y$  is called a regular value of  $\forall x \in f^{-1}(y)$  we have clf(x) is surjective  $\underbrace{AND}_{T}$  df(x) has a right inverse.

In finite dim' | setting surjectivity => = 1 right inverse

In a din' setting, I right imore off ter (df(x)) has a complement in Tox.

In particular, when f is Fredholm, when yo Y is a regular value of  $f: X \to Y$  if df(x) is surjective (enough!)

 $\Rightarrow$  ind(f) = dim(Kerdf(x)) for  $\forall x \in f^{-1}(y)$ .

Thu (Implicit function Thus).

 $f: X \longrightarrow Y$ ,  $y \in Y$  a regular value of f, then  $M:=f^{-1}(y)(c X)$  is a Banach wild and  $T_{*}M = \ker \operatorname{alf}(x) \ \forall x \in M$ .

Moreover, if f is Fredholm then dim M = ind(f) (fruite!).

Thun (Smale) Let X, Y separable Barach until, f. X > Y (smurch)

Fredholm map. Then { regular value ye Y } is residual, i.e.

contains a countable intersection of open and clease sets.

Back to our T-hol cure setting:

$$\mathcal{E} = \mathcal{O}_{l''}(\Sigma'')$$


$$\mathcal{E} = \mathcal{O}_{l''}(\Sigma'')$$

$$\mathcal{E} = \mathcal{O}_{l''}(\Sigma'')$$


of local comportation of Dr.

 $D_{u}(\zeta) = \overline{\partial_{J}} \zeta + \frac{1}{2} (\partial_{S} J)(u) \partial_{v} u d_{s} - (\partial_{K} J)(u) \partial_{s} u dt) \qquad (4)$ 

In terms of a local basis fer, ... ean? along u(small work of a pt in E)



here, either in (5,+)-coordinate or in (2, E) -coordinate



each  $f \in W^{kp}(u^{e_{TM}}|_{\Omega})$  is unitten by  $\int_{\Omega} = \int_{\Omega} e_{1} + \cdots + \int_{\Omega} e_{2n} \qquad f_{1} \cdot \Omega \to \mathbb{R}^{2n} \circ C^{n}$ and  $f \in W^{kp}(\Omega; C^{n})$ 

In (4), only the first term  $\overline{J}_3$  involves the derivatives of  $f_i$ .

Also, in terms of  $(Z, \overline{Z})$ -coordinate, we have  $\overline{J}_3 f = \overline{J}_2 f$ 

Duf = Def + A(2) of when A: Wk+,P(IZ, End(C"))

World

(as a unitary, each component is a
four in Wk+,P(IZ, End))

(=) 25 } = Dyf - A(=) }

 $\frac{\text{Thu (lnt)}}{\text{in SFT-3}} \| \| \| \|_{K,p} \leq C \left( \| \| \| \|_{p} + \| \| \| \| \|_{p} - A(2) \| \|_{K-1,p} \right) \\
\leq C \left( \| \| \| \|_{K-1,p} + \| \| \| \| \| \| \| \|_{Lp} + \| \| A(2) \| \| \|_{Lp} \right)$ 

· WER CON WEIR + WEIR is closed under multiplication, we get

(1A(2) 11/ Emp = C' (1) 11/ Emp

=> 11/51/KP = C, (11 DOZ1/K+ + 11/51/K+1)

- · Wk, p cpt Wkri, p + useful lemma, ve get Du is semi-Fredholm.
- · By a dual argument (consider Dut), one conclude that Du is Fredholm. Exe.

therefore, in order to prove M= | moduli space = 2 (0), we of T-60 | ceme | we actually arms at this pt by an informal argument near eller of SFT-2.

Ruk If so, then

din  $M_J$  = din ker (Du)  $\forall u \in \partial_J^{-1}(0)$ One ambiguity:  $\partial_J^{-1}(0)$  may not be upoch) connected!  $\partial_J^{-1}(0) = \partial_J^{-1}(0) \cup \partial_J^{-1}(0) \cup \cdots$ A. B. htp class
of im(u) in  $H_2(M_3(R))$ we way to dempose (possibly with asymptotic ends or boundary conds).

=> une consider MAJ (i.e. fix J but with further top constraints).

and dim MAJ should depend on class A (later lectures).

Morene, in each converted component, (ay  $\partial_{J,A}(0)$ ,

{Ut} {te [on] > Duy > ind (Du) is ind of the parch

→ a benefit to coloulate ind (Du) from some 'special" n ∈ 2, A(0).

knk (a delicate point). Does ind (Dn) (or more precisely ker (Dn) or coker (Dn)) depends on the regularity degree k ? NO

J is sourch  $\Rightarrow$  ue W<sup>lf</sup>( $\sum_{k}$ M)  $\cap$   $C^{l}$   $\Rightarrow$  thun above applies for any k>l.

assumption | assettime is infant smooth |  $\Rightarrow$  ( $\in$  Ker(Du)  $\subset$  W<sup>trf</sup> and ui>  $\sum_{k}$ Lool.